

## REFERENCES

### RUNNING

Albracht K and Arampatzis A. Exercise-induced changes in triceps surae tendon stiffness and muscle strength affect running economy in humans. *Eur J Appl Physiol* 2013. Epub

McBeth JM, Earl-Boehm JE, Cobb SC, et al. Hip muscle activity during 3 side-lying hip-strengthening exercises. Delgado TL, Kubera-Shelton E, Robb R. et al. Effects of foot strike on low back posture, shock attenuation, and comfort in running. *Med Sci Sports Exerc* 2012. Epub

Kong PW, Koh TM, Tan WC, Wang YS. Unmatched perception of speed when running overground and on a treadmill. *Gait Posture* 2012;36:46-8.

Williams DS 3rd, Green DH, Wurzinger B. Changes in lower extremity movement and power absorption during forefoot striking and barefoot running. *Int J Sports Phys Ther* 2012;7:525-32. ses in distance runners. *J Athl Train* 2012; 47:15-23.

Wouters I, Almonroeder T, DeJariais B et al. Effects of a movement training program on hip and knee joint frontal plane running mechanics. *IJSPT* 2012; 7:637-646.

Ford K, Taylor-Hass J, Genthe K et al. Relationship between hip strength and trunk motion in college cross-country runners. *Med Sci Sports Exerc* 2012; Epub

Kulmala JP, Avela J, Pasanen K, Parkkari J. Forefoot Strikers exhibit lower running-induced knee loading than forefoot strikers. *Med Sci Sports Exer* 2013 June 6. Epub

Riley PO, Dicharry J, Frnaz J et al. A kinematics and kinetic comparison of overground and treadmill running. *Med Sci Sports Exerc* 2008;40:1093-1100,

Warne JP, Warrington GD . Four-week habituation to simulated barefoot running improves running economy when compared with shod running. *Scand J Med Sci Sports* 2012. Epub

Hayes P, Caplan N. Foot strike patterns and ground contact times during high-caliber middle-distance races. *J Sports Sci* 2012;30:1275-83.

Hatala KG, Dingwall HL, Wunderlich RE, Richmond BG. Variation in Foot Strike Patterns during Running among Habitually Barefoot Populations. *PLoS One* 2013;8. Epub Jan.

Nielsen RO, Nohr EA, et al. Classifying running-related injuries based upon etiology, with emphasis on volume and pace. *Int J Sports Phys Ther.* 2013 Apr;8(2):172-9.

Rasmussen CH, Nielsen RO, et al. Weekly running volume and risk of running-related injuries among marathon runners. *Int J Sports Phys Ther.* 2013 Apr;8(2):111-20.

Sheerin KR, Hume PA, et al. Effects of a lower limb functional exercise programme aimed at minimizing knee valgus angle on running kinematics in youth athletes. *Phys Ther Sport*. 2012 Nov;13(4):250-4. Epub.

Kendall KD, Patel C, et al. Steps toward the validation of the Trendelenburg test: the effect of experimentally reduced hip abductor muscle function on frontal plane mechanics. *Clin J Sport Med*. 2013 Jan;23(1):45-51.

Gehring D, Mornieux G, et al. Knee and Hip Joint Biomechanics are Gender-specific in Runners with High Running Mileage. *Int J Sports Med*. 2013 Jul 18. Epub.

Murphy K, Curry EJ, et al. Barefoot Running: Does it Prevent Injuries? *Sports Med*. 2013 Aug 30. Epub.

Lenhart RL, Thelen DG, et al. Increasing Running Step Rate Reduces Patellofemoral Joint Forces. *Med Sci Sports Exerc*. 2013 Aug 12. Epub.

van Poppel D, Scholten-Peeters GG, et al. Prevalence, incidence and course of lower extremity injuries in runners during a 12-month follow-up period. *Scand J Med Sci Sports*. 2013 Aug 19. Epub.

Tsatalas T, Giakas G, et al. The effects of eccentric exercise-induced muscle damage on running kinematics at different speeds. *J Sports Sci*. 2013;31(3):288-98. Epub.

Giandolini M, Arnal PJ, et al. Impact reduction during running: efficiency of simple acute interventions in recreational runners. *Eur J Appl Physiol*. 2013 Mar;113(3):599-609. Epub.

Lorenz DS, Pontillo M. Is There Evidence to Support a Forefoot Strike Pattern in Barefoot Runners? A Review. *Sports Health*. 2012 Nov-Dec;4(6):480-484.

Brindle RA, Milner CE, Zhang S, Fitzhugh EC. Changing step width alters lower extremity biomechanics during running. *Gait Posture* 2013 July 2. Epub.

Bonacci J, Vincenzino B, Spratford W, Collins P. Take your shoes off to reduce patellofemoral joint stress during running. *Br J Sports Med* 2013 July 13. Epub

Willy RW, Davis IS. Kinematic and Kinetic Comparison of Running in Standard and Minimalist Shoes. *Med Sci Sports Exerc* 2013. July 19. Epub

Goldmann JP et al. The potential of toe flexor muscles to enhance performance. *J Sports Sci* 2013; 31(4): 424-33.

Willy RW, Scholz JP, David IS. Mirror gait retraining for the treatment of patellofemoral pain in female runners. *Clin Biomech* 2012; 27(10): 1045-51.

Rodrigues P, Tenbroek T, Van Emmerik R, Hamill J. Evaluating runners with and without anterior knee pain using the time to contact the ankle joint complexes' range of motion boundary. *Gait Posture* 2013 June 25. Epub

## **ILIOTIBIAL BAND SYNDROME**

Meardon SA, Campbell S, Derrick TR. Step width alters iliotibial band strain during running. *Sports Biomech* 2012; 11:464-72.

Selkowitz DM, Beneck G, Powers C. Which Exercises Target the Gluteal Muscles While Minimizing Activation of the Tensor Fascia Lata? Electromyographic Assessment Using Fine-Wire Electrodes. *JOSPT* 2012.

van der Worp MP, van der Horst N, de Wijer A et al. Iliotibial band syndrome in runners: a systematic review. *Sports Med* 2012 Nov 1; 42(11): 969-92.

## **ORTHOTICS**

Boldt AR, Wilson JD, Barrios JA, Kernozek TW. Effects of medially wedged foot orthoses on knee and hip joint running mechanics in females with and without patellofemoral pain syndrome. *J Appl Biomech* 2012 July 5. Epub.

## **RUNNING SHOES**

Willwacher S, Konig M, Potthast W et al. Does specific footwear facilitate energy storage and return at the metatarso-phalangeal joint in running? *J Appl Biomech* 2012. Epub

Fong Yan A, Sinclair PJ, Hiller C et al. Impact attenuation during weight bearing activities in barefoot vs. shod conditions: A systematic review. *Gait Posture* 2012. Epub

Ridge S, Johnson A, Mitchell U, Hunter I et al. Foot bone marrow edema after 10-week transition to minimalist running shoes. *Med Sci Sports Exerc.* 2013 Feb 22. Epub

Bonacci J, Saunders PU, Hicks A, et al. Running in a minimalist and lightweight shoe is not the same as running barefoot: A biomechanical study. *Br J Sports Med* 2013. Epub Jan.

Giandolini M, Horais N, et al. Impact reduction through long-term intervention in recreational runners: midfoot strike pattern versus low-drop/low-heel height footwear. *Eur J Appl Physiol.* 2013 Apr 14. Epub.

Sobhani S, Bredeweg S, et al. Rocker shoe, minimalis shoe, and standard running shoe: A comparison of running economy. *J Sci Med Sport*, 2013 May 24. Epub.

Paisis P, Hanley B, et al. Cypriot and greek army military boot cushioning: ground reaction forces and subjective responses. *Mil Med*. 2013 Apr;178(4):e493-7.

Willson JD, Bjorhus JS, et al. Short term changes in running mechanics and foot strike pattern following introduction to minimalistic footwear. *PM R*. 2013 Aug 30. Epub.

Lilley K, Stiles V, et al. The influence of motion control shoes on the running gait of mature and young females. *Gait Posture*. 2013 Mar;37(3):331-5. Epub.

Willy RW, Davis IS. Kinematic and Kinetic Comparison of Running in Standard and Minimalist Shoes. *Med Sci Sports Exerc*. 2013 Jul 19. Epub ahead of print

Nielsen RO, et al. Foot pronation is not associated with increased injury risk in novice runners wearing a neutral shoe: a 1-year prospective cohort study. *Br J Sports Med* 2013. June 13. Epub

## **PATELLOFEMORAL PAIN SYNDROME**

Noehren B, Hamill J, Davis I. Prospective Evidence for a Hip Etiology in patellofemoral pain. *Med Sci Sports Exerc*. 2012 Dec 27. Epub.

Power CM, Bolga L et al. Patellofemoral Pain: Proximal, Distal and Local Factors; 2<sup>nd</sup> International Research Retreat. *J Orthop Sports Phys Ther* 2012: 42(6)

Barton CJ, Lack S, Malliaras P, Morrissey D. Gluteal muscle activity and patellofemoral pain syndrome: a systematic review. *Br J Sports Med* 2012 Sept 3. Epub.

Willson JD, Petrowitz I, Butler RJ, Kernozek TW. Male and female gluteal muscle activity and lower extremity kinematics during running. *Clinical Biomech* 2012 Sept 1. Epub.

Willy RW, Manal KT, Witvrov EE, Davis IS. Are mechanics different between male and female runners with patellofemoral pain? *Med Sci Sports Exerc* 2012 July 26. Epub.

Noehren B, Sanchez Z, Cunningham T, McKeon PO. The effect of pain on hip and knee mechanics during running in females with chronic patellofemoral pain. *Gait Posture* 2012 July;36(3): 596-9.

Bazett-Jones D, Cobb S, Huddleston W et al. Effect of patellofemoral pain on strength and mechanics following an exhaustive run. *Med Sci Sports Exerc*. 2013 Jan 30. Epub.

Bonacci J, Vicenzino B, et al. Take your shoes off to reduce patellofemoral joint stress during running. *Br J Sports Med*. 2013 Jul 13. Epub.

Lankhorst NE, Bierma-Zeinstra SM, et al. Factors associated with patellofemoral pain syndrome: a systematic review. *Br J Sports Med*. 2013 Mar;47(4):193-206. Epub.

## **ACHILLES TENDON**

Silbernagel KG, Willy R, Davis I. Pre injury and post injury running analysis along with measurements of strength and tendon length in a patient with a surgically repaired Achilles tendon rupture. *J Orthop Sports Phys Ther* 2012 June;42(6) 521-9.

Peltonan J, Cronin NJ, Stenroth L et al. Achilles tendon stiffness is unchanged one hour after a marathon. *J Exp Biol* 2012 Oct 15;215(pt20):3665-71.

Wyndow N, Cowan SM, et al. Triceps surae activation is altered in male runners with Achilles tendinopathy. *J Electromyogr Kinesiol.* 2013 Feb;23(1):166-72. Epub.

## **HAMSTRINGS**

Kilcoyne KG, Dickens JF, Keblish D et al. Outcomes of Grade I and Grade II hamstring injuries in intercollegiate athletes: a novel rehabilitation protocol. *Sports Health* 2011 Nov, 3(6): 528-33.

Fyfe J, Opar D, Williams M, Shield A. The role of neuromuscular inhibition on hamstring strain injury recurrence. *Electromyogr Kinesiol.* 2013 Feb 8. Epub.

## **MEDIAL TIBIAL STRESS SYNDROME, COMPARTMENT SYNDROME AND STRESS FRACTURE**

Nielson RO, Cederholm P, Buist I et al. Can GPS be used to detect deleterious progression in training volume among runners? *J Strength Cond Res* 2012 Sept 17. Epub.

Yagi S, Muneta T, Sekiya I. Incidence and risk factors for medial tibial stress syndrome and tibial stress fracture in high school runners. *Knee Surg Sports Traumatol Arthrosc* 2012 Aug 9. Epub.

Nazem, TG, Ackerman KE. The Female Athlete Triad. *Sports Health.* July Aug 2012.

Daoud AI, Geissler GJ, Wang F et al. Foot strike and injury rates in endurance runners: a retrospective study. *Med Sci Sports Exerc* 2012 July 44(7) 1325-34.

Diebal A, Gregory R, Alitz C, Gerber P. Forefoot running improves pain and disability associated with chronic exertional compartment syndrome. *Am J Sports Med* 2012 Vol 40(5) 1060-67.

Carson DW, Myer GD, Hewett TE et al. Increased plantar force and impulse in American football players with high arch compared to normal arch. *Foot* 2012 Nov 7. Epub

Wentz L, Lin PY, Ilich JZ, Haymen EM. Dietary and training predictors of stress fractures in female runners. *Int J Sport Nutr Exerc Metab* 2012 Oct: 22(5): 374-82.

Barrack MT, Ackerman KE, et al. Update on the female athlete triad. *Curr Rev Musculoskelet Med*. 2013 Apr 24. Epub.

Tenforde AS, Sayres LC, et al. Identifying Sex-Specific Risk Factors for Stress Fractures in Adolescent Runners. *Med Sci Sports Exerc*. 2013 Apr 11. Epub.

Reinking MF, Hayes AM, et al. The effect of foot orthotic use on exercise related leg pain in cross country athletes. *Phys Ther Sport*. 2012 Nov;13(4):214-8. Epub.

Liem BC, Truswell HJ, et al. Rehabilitation and return to running after lower limb stress fractures. *Curr Sports Med Rep*. 2013 May-Jun;12(3):200-7.

Cauthon DJ, Langer P, et al. Minimalist shoe injuries: Three case reports. *Foot (Edinb)*. 2013 May 10. Epub.

## **EVALUATION**

Lisman P, O'Connor FG, Duester PA, Knapik JJ. Functional Movement Screen and Aerobic fitness predict injuries in military training. *Med Sci Sports Exerc* 2012 Nov 2012. Epub.